

Vacuum Polarization of a Charged Massless Scalar Field on Cosmic String Spacetime in the Presence of a Magnetic Field

J. Spinelly ^{*}and E. R. Bezerra de Mello [†]

Departamento de Física-CCEN
Universidade Federal da Paraíba
58.059-970, J. Pessoa, PB
C. Postal 5.008
Brazil

February 1, 2008

Abstract

In this paper we consider a charged massless scalar quantum field operator in the spacetime of an idealized cosmic string, i.e., an infinitely long, straight and static cosmic string, which presents a magnetic field confined in a cylindrical tube of finite radius. Three distinct situations are taking into account in this analysis: *i*) a homogeneous field inside the tube, *ii*) a magnetic field proportional to $1/r$ and *iii*) a cylindrical shell with δ -function. In these three cases the axis of the infinitely long tube of radius R coincides with the cosmic string. In order to study the vacuum polarization effects outside the tube, we explicitly calculate the Euclidean Green function associated with this system for the three above situations, considering points in the region outside the tube. Having these Green functions we calculate the renormalized vacuum expectation values, $\langle \hat{\Phi}^*(x)\hat{\Phi}(x) \rangle_{Ren.}$ and $\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.}$,

^{*}E-mail: spinelly@fisica.ufpb.br

[†]E-mail: emello@fisica.ufpb.br

associated with the charged field. In the evaluation of these vacuum polarization effects, two contributions appear for the three models. The firsts are the standard ones due to the conical geometry of the spacetime and the magnetic flux. The second contributions appear as an extra term. They are corrections due to the finite thickness of the radius of the tube. These extra terms provide relevant contributions, even for points very far away from the system, like a long-range effect. PACS numbers: 98.80.Cq, 11.10.Gh, 11.27.+d

1. Introduction

One of the firsts models which present topological defects was investigated by Nielsen and Olesen [1] several years ago. In their analysis topologically stable solutions named vortices, appears as a consequence of spontaneous breakdown of local Abelian and non-Abelian gauge symmetries in a classical field theory. These solutions correspond to infinitely long objects which carry magnetic fluxes. Later, Garfinkle investigated the Abelian model considering its influence on the geometry of the spacetime [2]. He found, as in flat spacetime, static cylindrically symmetric solutions representing vortices. He also shown that asymptotically the spacetime around the cosmic string is a Minkowski one minus a wedge. Its core has a non-zero thickness, and the magnetic field vanishes outside of it. Two years later Linet [3] obtained, under some specific condition, exact solution for the metric tensor. He was able to show that the structure of the respective spacetime corresponds to a conical one, with the conicity parameter being expressed in terms of the energy per unity length of the vortex. The complete analysis about the behavior of the gauge and matter fields near the cosmic string's core can only be obtained numerically. Some recent numerical analysis [4] about the structure of supermassive cosmic strings show that two different kind of solutions for the metric tensor exist.

A complete analysis about the behavior of a charged field in the neighborhood of a $U(1)$ -gauge cosmic string, must take into account not only the influence of the geometry of the spacetime but, also, the influence of the magnetic field. Two distinct cases can be analysed: *i)* The first one is to consider the string as an idealized infinitely thin linear topological defect, having a magnetic field running along it. This case can be treated analytically. *ii)* The second approach is to consider the non-zero thickness for the string. Analytically this problem becomes intractable. An intermediate approach is what we shall adopt here. We shall use an approximated model considering the spacetime produced by the string as being conical everywhere, but having a magnetic field inside a tube of non-zero thickness surrounding it. In this way some improvement is introduced when compared with the ideal case. Moreover, we shall admit specifics spatial behaviors for the magnetic field inside the tube trying to approach the real situation.

The vacuum polarization effects due to a magnetic field confined in a tube of finite radius in Minkowski spacetime has been first analysed by Sere-

bryanyi [5]. A few years later this effect in a charged massless scalar field on a idealized cosmic string spacetime, has been calculated by Guimarães and Linet [6]. There a magnetic flux running through the line singularity was considered. As a consequence the renormalized vacuum expectation value associated with the energy-momentum tensor, $\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.}$, presents a contribution coming from the geometry of the spacetime and the magnetic flux. In order to obtain this result the respective Green function was determined. In a particular choice of gauge, this Green function depends explicitly on the factor γ , the fractional part of the ratio of the magnetic flux ϕ by the quantum flux, $\phi_0 = 2\pi/e$.

More recently Sriramkumar [7] has calculated the vacuum fluctuation of current and energy densities for a massless charged scalar field around an idealized cosmic string carrying a magnetic flux. His approach to obtain the Green function takes into account the presence of the vector potential in the differential operator, $D_\mu = \partial_\mu - ieA_\mu$, which presents the advantage to calculate the two-points function without imposing any additional boundary condition on the field. Also the vacuum polarization associated with massless scalar field in a infinitely thin cosmic string embedded in a Randall-Sundrum type 2 brane world, has been calculated [8]. In lowest order of perturbation, $\langle T_{\mu\nu} \rangle_{Ren.}$, consists of two parts. The first one depends only on the conicity parameter and the second is a correction which depends on the parameter k connected with the cosmological constant.

Allen, Kay and Ottewill [9] have analyzed the vacuum polarization effect of a massless scalar field on a realistic cosmic string spacetime, considering generically the effect of the string's core through the non-minimal coupling between the scalar field with the geometry: $\xi \mathcal{R}$, being $\mathcal{R}$ the Ricci scalar and ξ the coupling constant ¹. In this case the respective Green function presents two parts: The first one, named "regular" by the authors, coincides with the usual one obtained in the literature for an idealized cosmic string, see [14] for example. The second contribution comes from the nonzero core's radius of the string. Although the radius of the cosmic string's core is very small ², the effect of this finite thickness on the vacuum polarization of a massless scalar field, is a long-range effects. Moreover it is of the same order

¹In [9] was not considered a specific model for the inner structure of the string; however, it was considered be infinitely long, straight and static.

²For cosmic string formed by a typical grand unified theory is of order 10^{-30} cm.

of magnitude as contribution coming from the geometry up to distance from the cosmic string that exceed the radius of the observable Universe.

Motivated by this result we raised a question about the consequence on the vacuum polarization of a charged massless scalar field in the $U(1)$ -gauge cosmic string spacetime considering this string surrounded by a magnetic field confined in a tube of finite radius. As in the Allen *et al* paper, we expect some consequence of the finite thickness of the tube on the vacuum polarization effect. In order to investigate this subject we decided in this paper, for simplicity, to analyse the behavior of a quantum charged massless scalar field in a idealized cosmic string spacetime, considering the presence of an infinitely long tube of finite radius R of magnetic field. Three distinct cases were taken into account:

- i)* a homogeneous magnetic field inside the tube,
- ii)* a magnetic field proportional to $1/r$ inside and
- iii)* a magnetic field on a cylindrical shell.

In all these three cases the axis of the infinitely long and straight tube coincides with the cosmic string.

The idealized model for an infinitely long straight static cosmic string spacetime can be given in cylindrical coordinates by the line element below:

$$ds^2 = -dt^2 + dr^2 + \alpha^2 r^2 d\theta^2 + dz^2, \quad (1)$$

where α is a parameter smaller than unity which codifies the presence of a conical two-surface (r, θ) . In fact for a typical Grand Unified Theory, $\alpha = 1 - O(10^{-6})$. (Here we considered that this cosmic string is lying along the z -direction.)

A magnetic field along the z -direction can be represented by $\vec{H}(r) = H(r)\hat{z}$. Now assuming that the field has a finite range in the radial coordinate, we are particularly interested in the three models below:

$$i) \quad H(r) = \frac{\phi}{\alpha\pi R^2}\Theta(R-r), \quad \text{homogeneous field inside} \quad (2)$$

$$ii) \quad H(r) = \frac{\phi}{2\pi\alpha Rr}\Theta(R-r), \quad \text{field proportional to } 1/r \text{ inside} \quad (3)$$

$$iii) \quad H(r) = \frac{\phi}{2\pi\alpha R}\delta(r-R), \quad \text{cylindrical shell}, \quad (4)$$

where R is the radial extent of the tube, and ϕ is the total flux. The ratio of

the flux by the quantum flux ϕ_o , can be expressed by $\phi/\phi_0 = N + \gamma$, where N is the integer part and $0 < \gamma < 1$.³

Considering both effects, due to the geometry of the spacetime and the finite thickness of the tube of magnetic field on the vacuum polarization of a charged massless scalar field, we observe that there appear two distinct contributions for each case: For the cases (i) and (ii), the firsts ones are due to the geometry and the magnetic flux. They are similar to the result found by Guimarães and Linet [6]. The seconds contributions are due the nonzero radius of the tube, R . The latter present similar dependence with the radial coordinate. In fact their dependence on the ratio R/r is of power type besides an overall $1/r^4$ dependence. These results indicate that the detailed information about the structure of the magnetic field inside the tube is not so relevant. The third case is completely different from the others. The second contribution to the vacuum polarization is now very similar to the result obtained in [9]. It presents a logarithmic dependence on the radial coordinate, being a long-range contribution.

Taking into account that the topological defects are of evident physical interest (they are considered as good candidates to explain some components of anisotropy on the cosmic microwave background [10], gamma ray burst [11], gravitational waves [12] and highest energy cosmic ray [13]) it seems reasonable to investigate the above mentioned problem in more detail.

This paper is organized as follows. In the section 2, we explicitly calculate the Green functions associated with this system for the three different models. Having now these Green functions, in section 3 we explicitly calculate the renormalized vacuum expectation values, $\langle \hat{\Phi}^*(x)\hat{\Phi}(x) \rangle_{Ren.}$, in complete form. Using a judicious criteria of approximation we present the behavior for these quantities. In section 4, we calculate the renormalized vacuum expectation values of the energy-momentum operator, $\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.}$. Using the same procedure as in the previous section we present approximated expressions for these results. We leave for section 5 our conclusions and most relevant remarks about this paper. Finally we want to say that we shall use in this paper $\hbar = G = c = 1$.

³These three models for the magnetic field has been considered previously by Bordag and Voropaev [15] to analyze the quantum mechanical behavior of spin-1/2 and gyro-magnetic $g \neq 2$ particle in a field of a magnetic string.

2. Green Function

The Green function associated with a charged massless scalar field must obey the non-homogeneous second order differential equation below

$$\frac{1}{\sqrt{-g}}\mathcal{D}_\mu \left[\sqrt{-g}g^{\mu\nu}\mathcal{D}_\nu \right] G(x, x') = -\delta^{(4)}(x - x'), \quad (5)$$

where $\mathcal{D}_\mu = \partial_\mu - ieA_\mu$.

Because it is easier to calculate the vacuum average of the square of the field and the energy-momentum tensor operators using the Euclidean version of the Green function, let us solve the above equation adopting the Euclidean extension of the cosmic string metric tensor given in (1), by performing a Wick rotation $t \rightarrow i\tau$. Now if we want to represent a magnetic field along the z -direction, the four-vector potential reads

$$A_\mu = (0, 0, A(r), 0) . \quad (6)$$

So the equation (5) becomes

$$\left[\frac{\partial^2}{\partial \tau^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{\partial^2}{\partial z^2} + \frac{1}{\alpha^2 r^2} \left(\frac{\partial^2}{\partial \theta^2} - 2ieA \frac{\partial}{\partial \theta} - e^2 A^2 \right) \right] G(x, x') = -\frac{1}{\alpha r} \delta(\tau - \tau') \delta(r - r') \delta(\theta - \theta') \delta(z - z') . \quad (7)$$

Now, in order to reproduce the magnetic fields configurations given by (2)-(4), we have only to write the third component of the vector potential by

$$A = \frac{\phi}{2\pi\alpha} a(r) . \quad (8)$$

For the two firsts models, we can represent the radial function $a(r)$ by:

$$a(r) = f(r)\Theta(R - r) + \Theta(r - R) , \quad (9)$$

with

$$f(r) = \begin{cases} r^2/R^2, & \text{for the model (i) and} \\ r/R, & \text{for the model (ii).} \end{cases} \quad (10)$$

For the third model,

$$a(r) = \Theta(R - r). \quad (11)$$

Due to the cylindrical symmetry of this system the Euclidean Green function can be expressed by

$$G(x, x') = \frac{1}{(2\pi)^3} \sum_{n=-\infty}^{\infty} e^{in(\theta-\theta')} \int_{-\infty}^{\infty} dk e^{ik(z-z')} \int_{-\infty}^{\infty} d\omega e^{i\omega(\tau-\tau')} g_n(r, r') . \quad (12)$$

Before to specialize on the specific model let us write down the non-homogeneous differential equation obeyed by the unknown function $g_n(r, r')$. Substituting (12) into (7) and using the standard representation to the delta functions in the temporal, angular and z -coordinates, we arrive to following differential equation for the function $g_n(r, r')$:

$$\left[\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{1}{\alpha^2 r^2} (n^2 - 2neA + e^2 A^2) - \beta^2 \right] g_n(r, r') = -\frac{1}{\alpha r} \delta(r - r') , \quad (13)$$

where $\beta^2 = k^2 + \omega^2$.

So, all the informations about the magnetic field, the radius of the cylindrical tube and the conicity of the spacetime are contained in $g_n(r, r')$. However, before to obtain specific solution, corresponding to specific model of magnetic field, some general properties must be satisfied by this function. Let us define by $g_n^<(r, r')$ the solution of (13) regular at $r \rightarrow 0$, and by $g_n^>(r, r')$ the solution that vanishes at infinity. These two solutions must satisfy the continuity condition at $r = r'$, with their first derivative discontinuous at this point.

It is of our main interest to investigate the vacuum polarization effect for external points to the magnetic flux. As we shall see, because the information about the field in this region must taking into account, the above mentioned effect will present contribution coming from the inner region. Now, let us consider solutions of (13) with both r and r' greater than R .

Let us first consider the models (i) and (ii). The inner solution, $g_n^<(r, r')$, of (13), corresponds to $r < r'$. However two regions must be considered separately: from 0 to R and from R to r' . Integrating out in region $r < R$, we have:

$$g_n^<(r, r') = \begin{cases} A_n^i H_i(r), & \text{for } r < R \\ B_n^i I_{|\nu|}(\beta r) + C_n^i K_{|\nu|}(\beta r), & \text{for } R < r < r', \end{cases} \quad (14)$$

where

$$\nu = \frac{(n - \delta)}{\alpha} , \quad (15)$$

being

$$\delta = \frac{e\phi}{2\pi\alpha} = N + \gamma . \quad (16)$$

$H_i(r)$, for $i = 1, 2$, represents the solution associated with the two firsts models:

$$H_1(r) = \frac{1}{r} M_{\sigma_1, \lambda_1} \left(\frac{\delta}{\alpha R^2} r^2 \right), \quad (17)$$

with $\sigma_1 = (\frac{n}{\alpha} - \frac{\beta^2 R^2 \alpha}{2\delta})/2$ and $\lambda_1 = n/2\alpha$, and

$$H_2(r) = \frac{1}{\sqrt{r}} M_{\sigma_2, \lambda_2}(\zeta r), \quad (18)$$

with $\sigma_2 = \frac{n\delta}{\alpha}(\delta^2 - \beta^2 R^2 \alpha^2)^{-1/2}$, $\lambda_2 = n/\alpha$ and $\zeta = \frac{2}{R\alpha}(\delta^2 + \beta^2 R^2 \alpha^2)^{1/2}$.

In both definitions above, Eq.s (17) and (18), $M_{\sigma, \lambda}$, represents the Whittaker function [16]. The continuity condition of $g_n^<$ and its first derivative at $r = R$, allow us to obtain the ratio $D_n^i = C_n^i/B_n^i$ as a function of βR only:

$$D_n^i(\beta R) = \frac{H'_i(R)I_{|\nu|}(\beta R) - H_i(R)I'_{|\nu|}(\beta R)}{H_i(R)K'_{|\nu|}(\beta R) - H'_i(R)K_{|\nu|}(\beta R)}. \quad (19)$$

In all the expressions, $I_{|\nu|}$ and $K_{|\nu|}$ are the modified Bessel functions.

The outer solution of (13) is given by

$$g_n^>(r, r') = E_n^i K_{|\nu|}(\beta r), \quad \text{for } r > r'. \quad (20)$$

All the above constants are determined by imposing the boundary conditions explained before on $g_n^<$ and $g_n^>$ at $r = r'$. In this way they will be function of r' . Finally we obtain the result to the complete radial Green function below:

$$g_n(r, r') = -\frac{1}{\alpha} K_{|\nu|}(\beta r_>) \left(I_{|\nu|}(\beta r_<) + D_n^i K_{|\nu|}(\beta r_<) \right) \quad \text{for } r \text{ and } r' > R. \quad (21)$$

In the above equation $r_>$ ($r_<$) is the bigger (smaller) value between r and r' . Substituting this solution into (12) and after some intermediate steps, the Euclidean Green function acquires the following expression:

$$G(x, x') = \frac{e^{iN\Delta\theta}}{8\pi^2 \alpha r r' \sinh u_0} \frac{e^{i\Delta\theta} \sinh(\gamma u_0/\alpha) + \sinh[(1-\gamma)u_0/\alpha]}{\cosh(u_0/\alpha) - \cos \Delta\theta}$$

$$\begin{aligned}
& + \frac{1}{4\pi^2} \int_0^\infty d\beta \beta J_0 \left(\beta \sqrt{(\Delta\tau)^2 + (\Delta z)^2} \right) \times \\
& \sum_{n=-\infty}^{\infty} e^{in\Delta\theta} D_n^i(\beta R) K_{|\nu|}(\beta r) K_{|\nu|}(\beta r'), \tag{22}
\end{aligned}$$

where

$$\cosh u_o = \frac{r^2 + r'^2 + (\Delta\tau)^2 + (\Delta z)^2}{2rr'}. \tag{23}$$

We can observe that the first term of the above Green function is, up to a gauge transformation, equivalent to the result presented by Guimarães and Linet [6] for the massless scalar field. The second term presents, besides the dependence on the α parameter and the magnetic flux, a dependence on the radius R of the tube through the constant D_n^i . Moreover, this Green function obeys the boundary condition $G(\theta' = \theta + 2\pi) = G(\theta' = \theta)$, which is a consequence of having used explicitly the vector potential in the differential equation (5). Although we have calculated this Green function for points outside the magnetic tube, there remains contributions coming from this region that do not disappear by taking $\gamma = 0$ and $\Delta\theta = 2\pi$.

As to the third model, the solution to $g_n(r, r')$ is:

$$g_n^<(r, r') = \begin{cases} A_n I_{|\nu|}(\beta r), & \text{for } r < R \\ B_n I_{|n/\alpha|}(\beta r) + C_n K_{|n/\alpha|}(\beta r), & \text{for } R < r < r', \end{cases} \tag{24}$$

and

$$g_n^>(r, r') = E_n K_{|n/\alpha|}(\beta r), \quad \text{for } r > r'. \tag{25}$$

Once more, the complete solution to this radial function can be obtained adopting a similar procedure as the previous case. This solution is:

$$g_n(r, r') = -\frac{1}{\alpha} K_{|n/\alpha|}(\beta r_>) \left(I_{|n/\alpha|}(\beta r_<) + D_n K_{|n/\alpha|}(\beta r_<) \right) \quad \text{for } r \text{ and } r' > R. \tag{26}$$

Again imposing the continuity condition on (24) and its first derivative at $r = R$ we get $D_n = C_n/B_n$ as function of βR :

$$D_n(\beta R) = \frac{I'_{|\nu|}(\beta R) I_{|n/\alpha|}(\beta R) - I_{|\nu|}(\beta R) I'_{|n/\alpha|}(\beta R)}{I_{|\nu|}(\beta R) K'_{|n/\alpha|}(\beta R) - I'_{|\nu|}(\beta R) K_{|n/\alpha|}(\beta R)}. \tag{27}$$

Finally adopting similar procedure as in the previous case, we obtain:

$$\begin{aligned}
G(x, x') &= \frac{1}{8\pi^2 \alpha r r' \sinh u_0} \left[\frac{\sinh(u_0/\alpha)}{\cosh(u_0/\alpha) - \cos \Delta\theta} \right] \\
&+ \frac{1}{4\pi^2} \int_0^\infty d\beta \beta J_0 \left(\beta \sqrt{(\Delta\tau)^2 + (\Delta z)^2} \right) \times \\
&\sum_{n=-\infty}^\infty e^{in\Delta\theta} D_n(\beta R) K_{|n|/\alpha}(\beta r) K_{|n|/\alpha}(\beta r') . \quad (28)
\end{aligned}$$

For this case we can see that the first term in this Green function coincides with results found by Smith [14] and by Linet in [17] for a massless scalar field. Moreover, the second term contains the information about the radius of the tube through the constant D_n .

So having the Green functions our next steps will be the calculations of the vacuum expectation values associated with the modulus of the scalar field and its energy-momentum tensor.

3. Computation of $\langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.}$

In this section we want to calculate the renormalized vacuum expectation values associated with the modulus square of the scalar field in this physical system for the three distinct configurations of magnetic fields. As we shall see these vacuum average will depend on the conicity parameter α , the fractional part of ϕ/ϕ_0 and the radius of the tube, R . Because the two firsts models present similar Green functions, we shall develop initially the calculations considering these two models together and later the third one.

The renormalized vacuum expectation value of the modulus square of the charged massless scalar field on idealized cosmic string spacetime is given by

$$\langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} = \lim_{x' \rightarrow x} [G(x, x') - G_E(x, x')], \quad (29)$$

where $G_E(x, x') = \frac{1}{4\pi^2(x-x')^2}$, is the Euclidean Green function for a flat four-dimensional spacetime, and $G(x, x')$ the Euclidean Green function associated with this system under investigation.

As we have explained before, we shall develop first this calculation for the models (i) and (ii), whose Green functions are given in (22) with D_n^i

given by (19). A very interesting aspect of these Green functions is that the correction due to the magnetic tube's radius is finite in the coincidence limit; so the result is:

$$\begin{aligned} \langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} &= \frac{1}{48\pi^2 r^2 \alpha^2} [1 - \alpha^2 - 6\gamma(1 - \gamma)] \\ &+ \frac{1}{(2\pi)^2} \int_0^\infty d\beta \beta \sum_{n=-\infty}^\infty D_n^i(\beta R) K_{|\nu|}^2(\beta r). \end{aligned} \quad (30)$$

In order to provide a quantitative information about the behavior of the correction in above equation, we must adopt the following procedure: *i*) Redefining a new variable $v = \beta r$, it is possible to see that the second term presents on overall $1/r^2$ dependence, the same as presented by first term. However there remains an extra radial dependence on the integrand through the coefficient $D_n^i(vR/r)$. So the complete information about the second term can only be obtained numerically specifying the ratio R/r . *ii*) An approximated behavior can be exhibited if we analyse the integrand of (30). Near the origin $K_{|\nu|}(v)$ behave as $v^{-2|\nu|}$, on the other hand $D_n^i(vR/r)$ present a dependence $v^{2|\nu|}$ as we shall see later. So the product $v D_n^i(vR/r) K_{|\nu|}^2(v)$ vanishes linearly with v . For large v , $v K_{|\nu|}^2(v)$ decays as e^{-2v} . Because $D_n^i(vR/r)$ grows slowly than $e^{2vR/r}$ for large v , its most significant relevance in the integrand happens only up v of order unity. However because we are mostly interested to study the vacuum polarization for points very far from the cosmic string, we shall consider $R/r \ll 1$ in the our analysis. So we may approximate the integrand of (30) assuming to the coefficient D_n^i its first order expansion. This is equivalent to consider the coefficients $D_n^i(\beta R)$ with $\beta R \ll 1$. Taking in (19) $\beta R \ll 1$, the most relevant term for both cases are given by

$$D_n^i(\beta R) = -\frac{2}{\Gamma(|\nu| + 1)\Gamma(|\nu|)} \left(\frac{g_i^n - |\nu| f_i^n}{g_i^n + |\nu| f_i^n} \right) \left(\frac{\beta R}{2} \right)^{2|\nu|}, \quad (31)$$

where

$$g_1^n = \left(\frac{\delta - n}{\alpha} - 1 \right) M_{\lambda_1, \lambda_2}(\delta/\alpha) + \left(1 + \frac{2n}{\alpha} \right) M_{\gamma_1, \lambda_1}(\delta/\alpha), \quad (32)$$

$$f_1^n = M_{\lambda_1, \lambda_2}(\delta/\alpha), \quad (33)$$

$$g_2^n = \left(\frac{\delta - n}{\alpha} - \frac{1}{2} \right) M_{\lambda_2, \lambda_1}(2\delta/\alpha) + \left(\frac{1}{2} + \frac{2n}{\alpha} \right) M_{\gamma_2, \lambda_2}(2\delta/\alpha) , \quad (34)$$

and

$$f_2^n = M_{\lambda_2, \lambda_1}(2\delta/\alpha) , \quad (35)$$

being $\gamma_1 = (n + 2\alpha)/2\alpha$, $\gamma_2 = (n + \alpha)/\alpha$ and $\nu = \frac{n-N-\gamma}{\alpha}$.

Using now these approximated expressions to the coefficients D_n^i , and with the help of [16], Eq. (30) becomes:

$$\begin{aligned} \langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} &= \frac{1}{48\pi^2 r^2 \alpha^2} [1 - \alpha^2 - 6\gamma(1 - \gamma)] \\ &- \frac{1}{(2\pi)^2 \alpha r^2} \sum_{n=-\infty}^{\infty} \frac{|\nu|}{1 + 2|\nu|} \left(\frac{g_i^n - |\nu| f_i^n}{g_i^n + |\nu| f_i^n} \right) \left(\frac{R}{r} \right)^{2|\nu|} \end{aligned} \quad (36)$$

The first term of the above result depends only on the fractional part of ϕ/ϕ_0 , besides the parameter α . On the other hand the second term, the correction, depends on the integer part of the the ratio ϕ/ϕ_0 , N . However, because R/r is assumed to be smaller than unity, the most relevant correction comes from the $n = N$. In this case the extra radial dependence of the second term is dominated by $(R/r)^{2\gamma/\alpha}$.

As we have already explained our approximation consists to substitute into (30), the expression (31) to the coefficients D_n^i . In fact doing this other contributions given by the complete expression (27) are missing; only the first order in the expansion was kept. In order to be sure that our approximation is a good one, we also analysed numerically the integrand of the second contribution of (30) using the exact expression (27) and the approximated result (31)-(35) for the coefficients for both cases $i = 1$ and 2 . For simplicity only we take in the summation $N = 0$, for both cases. This approximation is once more justified by the figures 1(a) and 1(b) which exhibit comparatively the integrand of (30) written in terms os the new variable v , $I(v) = v \sum D_n^i(vR/r) K_{|\nu|}^2(v)$, for the models (i) and (ii), respectively, considering the complete expressions for the coefficients and their respective approximations. We can see a good agreement between both curves for each case. (In these numerical analysis we have considered only the term $n = 0$ in the summation for both models. Considering other terms, for example n running from -20 to 20 , we have not observed any significant difference when

compared with the $n = 0$ term. So the summation in (36) can be mainly represented by its most relevant contribution.)

So accepting this fact as general, our final result to the vacuum polarization effects becomes:

$$\begin{aligned} \langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} &= \frac{1}{48\pi^2 r^2 \alpha^2} \left\{ \frac{1 - \alpha^2 - 6\gamma(1 - \gamma)}{\alpha^2} \right. \\ &\quad \left. - \frac{12\gamma}{\alpha(2\gamma + \alpha)} \left[\frac{g_i^N - \gamma/\alpha f_i^N}{g_i^N + \gamma/\alpha f_i^N} \right] \left(\frac{R}{r} \right)^{2\gamma/\alpha} \right\}. \end{aligned} \quad (37)$$

From the above expression, it is possible see that the second term has the same order of magnitude as the standard one, up to ratio $r/R = |\bar{x}|^{\alpha/2\gamma}$, being $\bar{x}$ given by

$$\bar{x} = \left[\frac{\alpha}{1 - \alpha^2 + 6\gamma(\gamma - 1)} \frac{12\gamma}{\alpha + 2\gamma} \left(\frac{g_i^N - \gamma/\alpha f_i^N}{g_i^N + \gamma/\alpha f_i^N} \right) \right]. \quad (38)$$

For smaller ratio the second contribution becomes more relevant than the first one. Once more, by numerical analysis we observe that for $N = 0$, $\alpha = 0.99$ and $\gamma = 0.3$, there happen a cancellation between both contributions for r/R of order 5 for both models. For $N = 1$ and assuming the same values for α and γ , the seconds terms changes their signs and there is no cancellation anymore.

After this discussion about the two firsts models, let us analyze the third one where the respective Green function is given by (28). For this case the renormalized vacuum expectation value of the modulus square of the scalar field becomes:

$$\begin{aligned} \langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} &= \frac{1}{48\pi^2 r^2 \alpha^2} (1 - \alpha^2) \\ &\quad + \frac{1}{(2\pi)^2} \int_0^\infty d\beta \beta \sum_{n=-\infty}^\infty D_n(\beta R) K_{|n|/\alpha}^2(\beta r), \end{aligned} \quad (39)$$

with the coefficients $D_n(\beta R)$ given by (27). Although the physical system that we are analysing is completely different from the one analyzed by Bruce Allen *et al* in [9], there appears a strong similarity between our result with theirs. Again if we are inclined to adopt an approximate expression to the

coefficients D_n considering $\beta R \ll 0$, two distinct behavior are obtained:
For $n \neq 0$,

$$D_n(\beta R) = -\frac{2}{\Gamma(|n|/\alpha + 1)\Gamma(|n|/\alpha)} \left(\frac{|\nu| - |n|/\alpha}{|\nu| + |n|/\alpha} \right) \left(\frac{\beta R}{2} \right)^{2|n|/\alpha}, \quad (40)$$

and for $n = 0$, and

$$D_0(\beta R) = \frac{\delta/\alpha}{\delta/\alpha \left[\ln \left(\frac{\beta R}{2} \right) + \mathcal{C} \right] - 1}, \quad (41)$$

where $\mathcal{C}$ is the Euler constant. As we can see for $n \neq 0$, $D_n(\beta R)$ vanishes at least as fast as $(\beta R)^{2|n|/\alpha}$ when $\beta R \rightarrow 0$. On the other hand, $D_0(\beta R)$ vanishes only with the inverse of the logarithm, so slowly. In this way the most relevant contribution to the summation in (39) comes from the $n = 0$. This same conclusion was previously obtained by Bruce Allen *et al* for another physical system. There they used a more convenient notation to express this coefficient, which also we adopt here:

$$D_0(\beta R) = \frac{1}{\ln(\beta/q)}, \quad (42)$$

where in our case

$$q = \frac{2}{R} e^{-\mathcal{C} + \alpha/\delta}. \quad (43)$$

So the most relevant contribution to the correction of the renormalized vacuum expectation value of the square of modulus of the scalar field in (39) is proportional to:

$$\int_0^\infty d\beta \beta \frac{K_0^2(\beta r)}{\ln(\beta/q)} = \frac{1}{r^2} \int_0^\infty dv v \frac{K_0^2(v)}{\ln(v) - \ln(qr)}. \quad (44)$$

As it was pointed out in [9] this approximation presents a problem because the above integrand has a pole at $v = qr$. However, this pole is a consequence the approximation adopted, and it occurs when the argument of D_0 is much greater than unity, where the approximation is no longer valid. Moreover analyzing the behavior of the integrand, they observe that $vK_0^2(v)$ vanishes at $v = 0$, peaks around v of order 1, and decays as e^{-2v} for large v . On the other hand D_0 vanishes for $v = 0$ and grows less slowly than $e^{2vR/r}$

for large v . So, finally they conclude that (44) can be well approximated discarding the term $\ln(v)$. (Also a numerical analysis in their paper supports this approximation.)

Here in our case a similar conclusion can be reached. In fact choosing $\gamma = 0.2$, $N = 0$ and $\alpha = 0.99$, $qr = \frac{2r}{R}e^{-\mathcal{C}+\alpha/\delta}$ in of order than 10^5 for $r/R = 10^3$, so $\beta R \approx 10^2$. Consequently the pole occurs in the region where the approximation is no longer valid. We shall not repeat their argument to accept the same approximation procedure; however we present in Fig. 2 a numerical evaluation of this approximation, comparing the integrand of the second contribution of (39) using the exact expression to its more relevant term, D_0 , given by (27) and its approximated expression given in (44), discarding the $\ln(v)$ term. So we can conclude, supported by those evidences, that the second term of (39) can be approximated by

$$-\frac{1}{(2\pi)^2} \frac{1}{r^2 \ln(qr)} \int_0^\infty dv v K_0^2(v). \quad (45)$$

Finally with the help of [16] we integrate the above equation, providing to (39) the expression below:

$$\langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} = \frac{1}{48\pi^2 r^2} \frac{1 - \alpha^2}{\alpha^2} - \frac{1}{8\pi^2 \alpha r^2 \ln(\frac{2r}{R} e^{-\mathcal{C}+\alpha/\delta})}. \quad (46)$$

From the above expression we can see that the explicit dependence on the flux tube appears only on the second term. Taking $R \rightarrow 0$ or $\delta \rightarrow 0$, this term vanishes. (The first term contains only the contribution coming from the geometry and vanishes when $\alpha \rightarrow 1$).

As in the physical system analyzed in [9], there is in this case a long-range effect in (46) coming from the special structure of the magnetic tube around the cosmic string.

Because $\alpha \approx 1$, (46) can be written as

$$\langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.} = \frac{(\alpha^{-1} - 1)}{24\pi^2 r^2} \left[1 - \frac{3}{(\alpha^{-1} - 1) \ln(\frac{2r}{R} e^{-\mathcal{C}+\alpha/\delta})} \right], \quad (47)$$

where we see that the second term inside the bracket vanishes slowly as $r \rightarrow \infty$. In fact it can be noticed that the second term is greater than the unity up to

$$\frac{r}{R} \approx \frac{1}{2} \exp \left[\frac{3}{\alpha^{-1} - 1} + \mathcal{C} - \frac{\alpha}{\delta} \right], \quad (48)$$

which $\alpha = 0.99$ and $\delta = 0.2$ we have $r/R = 10^{127}$. As in [9], even for $R = 10^{-30}cm$, the correction in (47) becomes more relevant than the geometrical effect up to $r \approx 10^{97}cm$ which exceed the radius of the observable Universe.

4. Computation of $\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.}$

The renormalized vacuum expectation value of the energy-momentum tensor associated with this system under investigation is given by:

$$\langle \hat{T}_\mu^\nu \rangle_{Ren.} = \lim_{x' \rightarrow x} [D_{\mu(\alpha, \phi)}^{\nu'} G(x, x') - D_{\mu(1,0)}^{\nu'} G_E(x, x')] , \quad (49)$$

with the operator $D_{\mu(\alpha, \phi)}^{\nu'}$ expressed as

$$D_{\mu(\alpha, \phi)}^{\nu'} = (1 - 2\xi) D_\mu \bar{D}^{\nu'} - \xi (D_\mu D^\nu + \bar{D}_\mu \bar{D}^{\nu'}) + (2\xi - \frac{1}{2}) \delta_\mu^\nu D_\sigma \bar{D}^{\sigma'} , \quad (50)$$

being $D_\sigma = \nabla_\sigma - ieA_\sigma$ and the bar denoting complex conjugate. Again, in order to take into account the presence of the three magnetic field configurations given in the beginning of this paper we write the four vector potential in the form $A_\sigma = (0, 0, \frac{\phi a(r)}{2\pi\alpha}, 0)$, with $a(r)$ being given by (9) and (10), for the firsts two cases and by (11) for the third one. The respective Green functions $G(x, x')$, are given by (22) and (28).

Because we are convince ourselves about the approximation procedure adopted in the last section, here we shall consider in all the calculations only the leading term of the coefficients D_n^i for each case.

Let us start with the two firsts models. After some intermediate calculations, we arrive at:

$$\begin{aligned} \langle \hat{T}_0^0(x) \rangle_{Ren.} &= \langle \hat{T}_0^0(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta D_{n=N}^i(\beta R) \left\{ 2\beta^2 \left(\xi - \frac{1}{4} \right) \times \right. \\ &\quad K_{\gamma/\alpha+1}^2(\beta r) + \left(2\xi\beta^2 + 4 \left(\xi - \frac{1}{4} \right) \right) \frac{\gamma^2}{\alpha^2 r^2} K_{\gamma/\alpha}^2(\beta r) \\ &\quad \left. - 4 \left(\xi - \frac{1}{4} \right) K_{\gamma/\alpha+1}(\beta r) K_{\gamma/\alpha}(\beta r) \right\} , \end{aligned} \quad (51)$$

$$\langle \hat{T}_1^1(x) \rangle_{Ren.} = \langle \hat{T}_1^1(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta D_{n=N}^i(\beta R) \left\{ \frac{\beta^2}{2} K_{\gamma/\alpha+1}^2(\beta r) \right.$$

$$\begin{aligned}
& - \left(2\xi + \frac{\gamma}{\alpha} \right) \frac{\beta}{r} K_{\gamma/\alpha+1}(\beta r) K_{\gamma/\alpha}(\beta r) \\
& - \left(\frac{\beta^2}{2} - \frac{2\xi\gamma}{\alpha r^2} \right) K_{\gamma/\alpha}^2(\beta r) \Big\} , \tag{52}
\end{aligned}$$

$$\begin{aligned}
\langle \hat{T}_2^2(x) \rangle_{Ren.} &= \langle \hat{T}_2^2(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta D_{n=N}^i(\beta R) \left\{ \left(2\beta^2\xi - \frac{\beta^2}{2} \right) \times \right. \\
& K_{\gamma/\alpha+1}^2(\beta r) + \left[2\xi - \frac{4\gamma}{\alpha} \left(\xi - \frac{1}{4} \right) \right] \frac{\beta}{r} K_{\gamma/\alpha+1}(\beta r) K_{\gamma/\alpha}(\beta r) \\
& + \left. \left[4\beta^2 \left(\xi - \frac{1}{4} \right) + \frac{4\xi\gamma}{\alpha r^2} \left(\frac{\gamma}{\alpha} - \frac{1}{2} \right) \right] K_{\gamma/\alpha}^2(\beta r) \right\} , \tag{53}
\end{aligned}$$

and by boost invariance in the $t - z$ plane $\langle \hat{T}_3^3(x) \rangle_{Ren.} = \langle \hat{T}_0^0(x) \rangle_{Ren.}$. Where the expression to $\langle \hat{T}_\mu^\nu(x) \rangle_{Reg.}$ is given by

$$\langle \hat{T}_\mu^\nu(x) \rangle_{Reg.} = u_1(\alpha, \gamma) diag(1, 1, -3, 1) + (\xi - 1/6) u_2(\alpha, \gamma) diag(1, -1/2, 3/2, 1) , \tag{54}$$

with

$$u_1(\alpha, \gamma) = \frac{1}{1440\pi^2 r^4} [(\alpha^{-4} - 1) - 30\alpha^{-4}\gamma(1 - \gamma^2)] \tag{55}$$

and

$$u_2(\alpha, \gamma) = \frac{1}{2\pi^2 r^4} \left[\frac{1}{6}(\alpha^{-2} - 1) + \alpha^{-2}\gamma(\gamma - 1) \right] . \tag{56}$$

Substituting the asymptotic expression for $D_{n=N}^i(\beta R)$, Eqs (31)- (35), into the above equations, after long calculation, we obtain:

$$\begin{aligned}
\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.} &= \langle \hat{T}_\mu^\nu(x) \rangle_{Reg.} + \frac{\gamma}{4\pi^2 r^4} \left(\frac{R}{r} \right)^{2\gamma/\alpha} \left(\frac{g_i^N - \gamma/\alpha f_i^N}{g_i^N + \gamma/\alpha f_i^N} \right) \times \\
& \frac{1}{\alpha^3(\alpha + 2\gamma)(3\alpha + 2\gamma)} [(\xi - 1/6) diag(v_1, v_2, v_3, v_1) \\
& + \frac{\gamma}{3} \left(\gamma - \frac{\alpha}{2} \right) diag(h_1, h_2, h_3, h_1)] , \tag{57}
\end{aligned}$$

where

$$v_1 = -(12\alpha^3 + 4\alpha\gamma(8\alpha + 7\gamma) + 8\gamma^3) , \tag{58}$$

$$v_2 = 6\alpha^3 + 2\alpha\gamma(5\alpha + 2\gamma) , \tag{59}$$

$$v_3 = -[6\alpha^3 + 2\alpha\gamma(5\alpha + 2\gamma)](3\alpha + 2\gamma) , \quad (60)$$

and

$$h_1 = 2\alpha + 2\gamma , \quad (61)$$

$$h_2 = 2\alpha , \quad (62)$$

$$h_3 = -6\alpha - 4\gamma . \quad (63)$$

From the above expressions we can see that: *i*) It reproduces the Guimarães and Linet's result in the limit $R \rightarrow 0$, *ii*) the corrections decrease with the power of (R/r) proportional to the fractional part of ϕ/ϕ_0 and *iii*) for $\xi = 1/6$ the trace of (57) vanishes.

Finally it should be mentioned that this expression satisfies the conservation condition

$$\nabla_\nu \langle \hat{T}_\mu^\nu(x) \rangle_{Ren.} = 0 . \quad (64)$$

As to the third model the respective Green function is given by (28) and the radial function $a(r)$ by (11). Adopting a similar procedure as in the last two previous calculations we obtain:

$$\begin{aligned} \langle \hat{T}_0^0(x) \rangle_{Ren.} &= \langle \hat{T}_0^0(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta^3 D_0(\beta R) \left\{ 2\xi K_0^2(\beta r) \right. \\ &\quad \left. + 2 \left(\xi - \frac{1}{4} \right) K_1^2(\beta r) \right\} , \end{aligned} \quad (65)$$

$$\begin{aligned} \langle \hat{T}_1^1(x) \rangle_{Ren.} &= \langle \hat{T}_1^1(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta^3 D_0(\beta R) \left\{ \frac{1}{2} K_1^2(\beta r) \right. \\ &\quad \left. - \frac{2\xi}{\beta r} K_0(\beta r) K_1(\beta r) - \frac{1}{2} K_0^2(\beta r) \right\} , \end{aligned} \quad (66)$$

$$\begin{aligned} \langle \hat{T}_2^2(x) \rangle_{Ren.} &= \langle \hat{T}_2^2(x) \rangle_{Reg.} + \frac{1}{4\pi^2\alpha} \int_0^\infty d\beta \beta^3 D_0(\beta R) \left\{ 2(\xi - 1/4) K_1^2(\beta r) \right. \\ &\quad \left. + \frac{2\xi}{\beta r} K_0(\beta r) K_1(\beta r) + 2(\xi - 1/4) K_0^2(\beta r) \right\} , \end{aligned} \quad (67)$$

with $\langle \hat{T}_3^3(x) \rangle_{Ren.} = \langle \hat{T}_0^0(x) \rangle_{Ren.}$. Now we have

$$\begin{aligned} \langle \hat{T}_\mu^\nu(x) \rangle_{Reg.} &= \frac{1}{1440\pi^2 r^4} (\alpha^{-4} - 1) diag(1, 1, -3, 1) \\ &\quad + \frac{1}{24\pi^2 r^4} (\alpha^{-2} - 1) (\xi - 1/6) diag(2, -1, 3, 2) , \end{aligned} \quad (68)$$

which is the result for the renormalized vacuum expectation value of the energy-momentum tensor associated with a massless scalar field on a idealized cosmic string spacetime [14, 17]. This result, together with (39), seem very curious, because although we are considering a charged field in the presence of a cylindrical shell of magnetic field, differently from the two previous cases, the electromagnetic effect appears only on the second terms for both calculations; moreover this effect is of long-range type.

Finally, substituting (42) into (65)-(67), taking into account the same considerations to overcome the integral problem on vacuum expectation value of the square of the scalar field, we obtain:

$$\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.} = \langle \hat{T}_\mu^\nu(x) \rangle_{Reg.} - \frac{1}{4\pi^2 \alpha r^4 \ln(qr)} (\xi - 1/6) \text{diag}(2, -1, 3, 2) . \quad (69)$$

It is easy to see that this result is traceless when $\xi = 1/6$ as conformal invariance demands on this spacetime; however as notice by Bruce Allen *et al* for $\xi \neq 1/6$, (69) presents a small violation of the conservation condition, which vanishes for $R \rightarrow 0$, or $q \rightarrow \infty$. Being $\xi \neq 1/6$, the energy density associated with a massless scalar charged field in this spacetime presents a long-range corrections arising from the magnetic field. For realistic models where the parameter α is very close to the unity this correction becomes even more relevant than the contribution coming from the geometry itself up to distance of the order the radius of the observable Universe.

5. Concluding Remarks

In this paper we have analyzed the vacuum polarization effects of a charged massless scalar field due to the locally flat cosmic string spacetime and a magnetic flux confined on a tube.

Three distinct configurations of magnetic flux have been considered. Among them, the cylindrical shell of δ -function field presents a long-range effect on the vacuum polarization.

In general, the results obtained to $\langle \hat{T}_\mu^\nu(x) \rangle_{Ren.}$ as to $\langle \hat{\Phi}^*(x) \hat{\Phi}(x) \rangle_{Ren.}$, outside the tube, present corrections due to the magnetic tube which vanish when we take $\phi \rightarrow 0$ or $R \rightarrow 0$.

Specifically speaking about the trace anomaly, it is well known that [18]

$$\langle \hat{T}_\mu^\mu(x) \rangle_{Ren.} = \frac{a_2(x)}{16\pi^2} , \quad (70)$$

where $a_2(x)$ presents contributions coming from the geometry of the space-time as well as contribution from the gauge field strength [19]. However because the spacetime under consideration is locally flat, and the region of our interest has no magnetic field we have $a_2(x) = 0$.

We conclude this paper by mentioning the importance to consider a finite structure of the magnetic flux on the quantum aspects of a charged field on an Abelian cosmic string spacetime. Although the complete knowledge about this structure could not be very well understood, its influence on the vacuum polarization effect can be even more relevant than the influence of geometry itself.

Acknowledgments

We would like to thank to V. M. Mostepanenko for a critical and enthusiastic reading of this paper. Also we would like to thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq.) and CAPES for partial financial support.

References


- [1] N. B. Nielsen and P. Olesen, Nucl. Phys. **B61**, 45 (1973).
- [2] D. Garfinkle, Phys. Rev. D **32**, 1323 (1985).
- [3] B. Linet, Phys. Lett. B **124**, 240 (1987).
- [4] M. Christensen, A. L. Larsen and Y. Verbin, Phys. Rev. D **60**, 125012 (1999); Y. Brihaye and M. Lubo, *ibid* **62**, 085005 (2000).
- [5] E. M. Serebryanyi, Theor. Math. Phys. **64**, 846 (1985).
- [6] M. E. X. Guimarães and B. Linet, Commun. Math. Phys. **165**, 297 (1994).
- [7] L. Sriramkumar, Class. Quantum Grav. **18**, 1015 (2001).
- [8] Y. Grats and A. Rossikhin, Modern Phys. Lett A **17**, 1207 (2002).
- [9] B. Allen, B. S. Kay and A. C. Ottewill, Phys. Rev. D **53**, 6829 (1996).
- [10] S. Sarangi and S. -H. Henry Tye, Phys. Lett. B **536**, 185 (2002), and K. Benabed and F. Bernardeu, Phys. Rev. D **62**, 123510 (2000).
- [11] V. Berezhinski, B. Hnatyk and A. Vilenkin, Phys. Rev. D **64**, 043004.
- [12] T. Damour and A. Vilenkin, Phys. Rev. Lett. **85**, 3761 (2000).
- [13] P. Bhattacharjee and G. Sigl, Phys. Rep. **327**, 109 (2000).
- [14] A. G. Smith, in *Symposium on the Formation and Evolution of Cosmic String*, edited by G. W. Gibbons, S. W. Hawking and T. Vachaspati (Cambridge University Press, Cambridge, England, 1989).

- [15] M. Bordag and S. Voropaev, J. Phys. A: Math. Gen. **26**, 7637 (1993).
- [16] I. S. Gradshteyn and I. M Ryzhik, *Table of Integral, Series and Products* (Academic Press, New York, 1980).
- [17] B. Linet, Phys. Rev. D, **35**, 536 (1987).
- [18] N. D. Birrell and P. C. W. Davies. *Quantum Fields in Curved Space* (Cambridge University Press. Cambridge. England. 1982)
- [19] I. Jack and L. Parker, Phys. Rev. D **31**, 2439 (1985).


6. Figure Captions

Figure 1: The solid curve in Fig. 1(a) represents the integrand of Eq. (30) written in terms of the new variable $v = \beta r$ for the first model considering the complete expression for the coefficient D_0^1 given by (19). The dashed curve is the same integrand now using the approximated expression for D_0^1 , Eq. (31). A similar analysis for the second model is given in Fig. 1(b). In both numerical analysis we have used $\alpha = 0.99$, $\gamma = 0.2$, $N = 0$ and $R/r = 10^{-3}$.

Figure 2: The solid curve in Fig. 2 represents the integrand of (39) for the third model considering the exact expression for D_0 , given by (27). The dashed curve exhibits the same integrand considering now the approximated expression to D_0 . In this numerical analysis we have used $\alpha = 0.99$, $\gamma = 0.2$, $N = 0$ and $R/r = 10^{-3}$.



(a)



(b)

Figure 1:




Figure 2: